

Popular Distributions

Shin-ichi Mayekawa

mayekawa@nifty.com

<http://mayekawa.in.coocan.jp/>

Notation

- notation

RV distributed as `dist_name(parameters)`

~

pdf or pf:

$$f(\textcolor{brown}{x} | \text{parameters}) = c \times k(x)$$

normalization constant

$$\int_X k(x) dx = \frac{1}{c}$$

Moments

$$mean = \mathcal{E}(X) = \int_X x f_X(x) dx$$

$$var(X) = \int_X (x - \mathcal{E}(X))^2 f_X(x) dx = \int_X x^2 f_X(x) dx - (\mathcal{E}(X))^2$$

$$\begin{aligned} cov(X_i, X_j) &= \int_{X_i} \int_{X_j} (x_i - \mathcal{E}(X_i))(x_j - \mathcal{E}(X_j)) f(x_i, x_j) dx_i dx_j \\ &= \int_{X_1} \int_{X_2} \cdots \int_{X_p} (x_i - \mathcal{E}(X_i))(x_j - \mathcal{E}(X_j)) f_X(x_1, x_2, \dots, x_p) dx_p \cdots dx_2 dx_1 \end{aligned}$$

$$\mathcal{D}(\mathbf{X}) = \begin{bmatrix} var(X_1) & cov(X_1, X_2) & \dots & cov(X_1, X_j) & \dots & cov(X_1, X_p) \\ cov(X_2, X_1) & var(X_2) & \dots & cov(X_2, X_j) & \dots & cov(X_2, X_p) \\ \vdots & \vdots & \dots & \dots & \dots & \vdots \\ cov(X_i, X_1) & cov(X_i, X_2) & \dots & cov(X_i, X_j) & \dots & cov(X_i, X_p) \\ \vdots & \vdots & \dots & \dots & \dots & \vdots \\ cov(X_p, X_1) & cov(X_p, X_2) & \dots & cov(X_p, X_j) & \dots & var(X_p) \end{bmatrix}$$

3

Moments

- calculation of the dispersion matrix

$$\mathcal{E}(\mathbf{X}) = (\mathcal{E}(X_1), \mathcal{E}(X_2), \dots, \mathcal{E}(X_p))'$$

$$\mathcal{D}(\mathbf{X}) = \mathcal{E}(\mathbf{X}\mathbf{X}') - \mathcal{E}(\mathbf{X})\mathcal{E}(\mathbf{X})'$$

- using conditional expectation and dispersion

$$\mathcal{E}(X_1) = \mathcal{E}(\mathcal{E}(X_1|X_2))$$

$$var(X_1) = \mathcal{E}(var(X_1|X_2)) + var(\mathcal{E}(X_1|X_2))$$

$$\mathcal{E}(X_1) = \mathcal{E}(\mathcal{E}(X_1|\mathbf{X}_2))$$

$$\mathcal{D}(X_1) = \mathcal{E}(\mathcal{D}(X_1|\mathbf{X}_2)) + \mathcal{D}(\mathcal{E}(X_1|\mathbf{X}_2))$$

4

Posterior Variance

$$var(X_1) = \mathcal{E}(var(X_1|X_2)) + var(\mathcal{E}(X_1|X_2))$$

$$X_1 \rightarrow \theta \quad X_2 \rightarrow X$$

$$\mathcal{E}_X[var(\theta|X)] = var(\theta) - var[\mathcal{E}(\theta|X)]$$

5

Mode of pdf

- mode = value of RV at which pdf is maximum.

$$\frac{d f_X(x)}{d x} = 0 \quad \text{or} \quad \frac{d \log f_X(x)}{d x} = 0$$

$$\frac{\partial f(x,y)}{\partial x} = 0 \quad \text{and} \quad \frac{\partial f(x,y)}{\partial y} = 0$$

6

Bernoulli

- notation

$$X \sim Bernoulli(\pi)$$

- pf

$$f(x|\pi) = \pi^x (1-\pi)^{1-x}$$

- moments

$$\mathcal{E}(X) = \pi$$

$$var(X) = \pi(1-\pi)$$

7

Binomial

- notation

$$X \sim Bin(n, \pi)$$

- pf

$$f(x|n, \pi) = {}_n C_x \times \pi^x (1-\pi)^{n-x} = \frac{n!}{x!(n-x)!} \times \pi^x (1-\pi)^{n-x}$$

- moments

$$\mathcal{E}(X) = n\pi \quad var(X) = n\pi(1-\pi)$$

$$mode(X) = [(n+1)\pi]$$

8

Multinomial

- notation $X \sim Multin(n, \boldsymbol{\pi})$
- pf $\sum_{j=1}^k x_j = n$ $\sum_{i=1}^k \pi_i = 1$

$$f(\mathbf{x}|n, \boldsymbol{\pi}) = \frac{n!}{x_1!x_2!\dots x_k!} \quad \pi_1^{x_1}\pi_2^{x_2}\dots\pi_k^{x_k} = \frac{n!}{\prod_{i=1}^k x_i!} \cdot \prod_{i=1}^k \pi_i^{x_i}$$

- moments

$$\mathcal{E}(X_i) = n\pi_i \quad var(X_i) = n\pi_i(1 - \pi_i)$$

9

Poisson

- notation $X \sim Poisson(\mu)$
- pf

$$f(x|\mu) = \exp(-\mu) \times \frac{\mu^x}{x!}$$

- moments

$$\mathcal{E}(X) = \mu \quad var(X) = \mu$$

$$mode = \mu - 1 \quad \text{if } \mu > 1$$

- comment

$$f(x|\mu) = \frac{n!}{x!(n-x)!} \left(\frac{\mu}{n}\right)^x \left(1 - \frac{\mu}{n}\right)^{n-x}$$

10

Normal

- notation $X \sim N(\mu, \sigma^2)$

- pdf

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \times \exp\left(-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}\right)$$

- moments

$$\mathcal{E}(X) = \mu \quad \text{var}(X) = \sigma^2$$

11

Multivariate Normal

- notation $X \sim MVN(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

- pdf

$$f(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{p/2}|\boldsymbol{\Sigma}|^{1/2}} \times \exp\left(-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})\right)$$

- example

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_1^2 & \sigma_{12} \\ \sigma_{12} & \sigma_2^2 \end{pmatrix} \quad \rho = \frac{\sigma_{12}}{\sigma_1\sigma_2}$$

$$\frac{1}{2\pi\sqrt{1-\rho^2}} \times \exp\left(-\frac{1}{2(1-\rho^2)} \left[\left(\frac{x-\mu_1}{\sigma_1}\right)^2 - 2\rho \left(\frac{x-\mu_1}{\sigma_1}\right) \left(\frac{y-\mu_2}{\sigma_2}\right) + \left(\frac{y-\mu_2}{\sigma_2}\right)^2 \right]\right)$$

- moments

$$\mathcal{E}(\mathbf{X}) = \boldsymbol{\mu} \quad \mathcal{D}(\mathbf{X}) = \boldsymbol{\Sigma}$$

12

Multivariate Normal (cont.)

- linear combinations

$$A\mathbf{X} + \mathbf{b} \sim N(\mathbf{b} + A\boldsymbol{\mu}, A\Sigma A')$$

- from conditional and marginal

$$\mathbf{y}_2 | \mathbf{y}_1 \sim MVN(\boldsymbol{\gamma}_2 + \mathbf{G}\mathbf{y}_1, \Psi)$$

$$\mathbf{y}_1 \sim MVN(\boldsymbol{\gamma}_1, \Phi)$$

$$\mathbf{y}_2 \sim MVN(\boldsymbol{\gamma}_2 + \mathbf{G}\boldsymbol{\gamma}_1, \mathbf{G}\Phi\mathbf{G}' + \Psi)$$

- distribution of the quadratic form

$$Q = \frac{1}{p}(\mathbf{x} - \boldsymbol{\mu})'\Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu}) \sim \chi^2(p, 1)$$

13

t

- notation $X \sim t(\nu, \gamma, c)$

- pdf

$$f(x|\nu, \gamma, c) = \frac{\Gamma(\frac{1}{2}(\nu+1))}{\Gamma(\frac{1}{2})\Gamma(\frac{\nu}{2})} \nu^{-\frac{1}{2}} c^{-\frac{1}{2}} \times \left(1 + \frac{1}{\nu c}(x - \gamma)^2\right)^{-\frac{1}{2}(\nu+1)}$$

- moments

$$\mathcal{E}(X) = \gamma \quad \text{if } \nu > 1$$

$$var(X) = \frac{\nu}{\nu - 2}c \quad \text{if } \nu > 2$$

- distribution of the quadratic form

$$Q = \frac{1}{c}(X - \gamma)^2 \sim F(1, \nu)$$

14

Multivariate t

- notation $X \sim MVt(\nu, \gamma, C)$

- pdf

$$f(x|\nu, \gamma, C) = \frac{\Gamma(\frac{1}{2}(\nu+p))}{[\Gamma(\frac{1}{2})]^p \Gamma(\frac{\nu}{2})} \nu^{-\frac{1}{2}p} |C|^{-\frac{1}{2}} \times \left(1 + \frac{1}{\nu}(x - \gamma)' C^{-1} (x - \gamma)\right)^{-\frac{1}{2}(\nu+p)}$$

- moments

$$\mathcal{E}(X) = \mu, \quad \mathcal{D}(X) = \frac{\nu}{\nu - 2} C$$

- distribution of the quadratic form

$$Q = \frac{1}{p}(x - \gamma)' C^{-1} (x - \gamma) \sim F(p, \nu)$$

15

Gamma

- notation $X \sim \Gamma(\alpha, \beta)$

- pdf

$$f(x|\alpha, \beta) = \frac{1}{\Gamma(\alpha)\beta^\alpha} \times x^{\alpha-1} \exp\left(-\frac{x}{\beta}\right)$$

$$\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} \exp(-x) dx$$

- moments

$$\mathcal{E}(X) = \alpha\beta \quad var(X) = \alpha\beta^2$$

$$mode(X) = \beta(\alpha - 1)$$

16

Chi Squared

- notation
- pdf

$$X \sim \Gamma\left(\frac{\nu}{2}, 2\right) \longleftrightarrow X \sim \chi^2(\nu)$$

$$f(x|\nu) = \frac{1}{\Gamma(\frac{\nu}{2})2^{\frac{\nu}{2}}} \times x^{\frac{\nu}{2}-1} \exp\left(-\frac{x}{2}\right)$$

- moments

$$\begin{aligned}\mathcal{E}(X) &= \nu\kappa^2 & \text{var}(X) &= 2\nu(\kappa^2)^2 \\ \text{mode}(X) &= \kappa^2(\nu - 2)\end{aligned}$$

- relationship

$$X/\kappa^2 \sim \chi^2(\nu), \quad \frac{1}{X} \sim \chi^{-2}(\nu, \frac{1}{\kappa^2}), \quad \sqrt{X} \sim \chi(\nu, \kappa^2)$$

17

Inverted Chi Squared

- notation
- pdf

$$f(x|\nu, \kappa^2) = \frac{(\kappa^2)^{\frac{\nu}{2}}}{2^{\nu/2}\Gamma(\frac{\nu}{2})} \times x^{-\frac{\nu}{2}-1} \exp\left(-\frac{\kappa^2}{2x}\right)$$

- moments

$$\begin{aligned}\mathcal{E}(X) &= \frac{\kappa^2}{\nu - 2} & \text{var}(X) &= \frac{2(\kappa^2)^2}{(\nu - 2)^2(\nu - 4)} \\ \text{mode}(X) &= \frac{\kappa^2}{\nu + 2}\end{aligned}$$

- relationship

$$\frac{\kappa^2}{X} \sim \chi^2(\nu), \quad \frac{1}{X} \sim \chi^2(\nu, \frac{1}{\kappa^2}), \quad \sqrt{X} \sim \chi^{-1}(\nu, \kappa^2)$$

18

Inverted Chi

- notation $X \sim \chi^{-1}(\nu, \kappa^2)$ $\kappa^2/X^2 \sim \chi^2(\nu)$
- pdf

$$f(x|\nu, \kappa^2) = \frac{(\kappa^2)^{\frac{\nu}{2}}}{2^{\frac{\nu}{2}-1}\Gamma(\frac{\nu}{2})} \times x^{-\nu-1} \exp\left(-\frac{\kappa^2}{2x^2}\right)$$

- moments

$$\begin{aligned} \mathcal{E}(X) &= \frac{\kappa\Gamma(\frac{\nu-1}{2})}{\sqrt{2}\Gamma(\frac{\nu}{2})} \approx \frac{\kappa}{\sqrt{\nu-\frac{3}{2}}} & \text{var}(X) &= \frac{\kappa^2}{\nu-2} - (\mathcal{E}(X))^2 \\ \text{mode}(X) &= \frac{\kappa}{\sqrt{\nu+1}} & & \approx \frac{\kappa^2}{2(\nu-2)(\nu-\frac{5}{3})} \end{aligned}$$

- relationship

$$\frac{\kappa^2}{X^2} \sim \chi^2(\nu), \quad X^2 \sim \chi^{-2}(\nu, \kappa^2), \quad \frac{1}{X} \sim \chi(\nu, \frac{1}{\kappa^2})$$

19

F

- notation $X \sim F(\nu_1, \nu_2)$ $\frac{1}{X} \sim F(\nu_2, \nu_1)$
- pdf

$$f(x|\nu_1, \nu_2) \sim \frac{\nu_1^{\nu_1/2} \nu_2^{\nu_2/2}}{B(\frac{1}{2}\nu_1, \frac{1}{2}\nu_2)} \frac{x^{\frac{\nu_1}{2}-1}}{(\nu_2 + \nu_1 x)^{(\nu_1+\nu_2)/2}}$$

- moments

$$\begin{aligned} \mathcal{E}(X) &= \frac{\nu_2}{\nu_2-2}, & \text{var}(X) &= \frac{2\nu_2^2(\nu_1+\nu_2-2)}{\nu_1(\nu_2-2)^2(\nu_2-4)} \\ \text{mode}(X) &= \frac{\nu_2}{\nu_2+1} \frac{\nu_1-2}{\nu_1} \end{aligned}$$

- relationship

$$\begin{aligned} x_1 &\sim \chi^{-2}(\nu_1, \kappa_1^2) \\ x_2 &\sim \chi^{-2}(\nu_2, \kappa_2^2) \end{aligned}$$

$$F = \frac{\frac{x_2}{\kappa_2^2} \times \frac{1}{\nu_2}}{\frac{x_1}{\kappa_1^2} \times \frac{1}{\nu_1}}$$

20

Normal-Inverted Chi

- notation

$$\begin{pmatrix} \mu \\ \sigma \end{pmatrix} \sim N\chi^{-1}(\gamma, \phi, \nu, \kappa^2)$$

- pdf

$$f(\mu, \sigma | \gamma, \phi, \nu, \kappa^2) = h(\mu | \sigma)h(\sigma)$$

$$\begin{aligned} &\propto \frac{1}{\sqrt{\sigma^2 \phi}} \exp\left(-\frac{1}{2\sigma^2}(\mu - \gamma)^2 \frac{1}{\phi}\right) \times \sigma^{-\nu-1} \exp\left(-\frac{\kappa^2}{2\sigma^2}\right) (\kappa^2)^{\frac{1}{\nu}} \\ &\propto \sigma^{-\nu-2} \exp\left[-\frac{1}{2\sigma^2} \left(\frac{1}{\phi}(\mu - \gamma)^2 + \kappa^2\right)\right] \end{aligned}$$

- definition

$$\mu | \sigma \sim N(\gamma, \sigma^2 \phi)$$

$$\sigma \sim \chi^{-1}(\nu, \kappa^2)$$

$$\mu \sim t(\nu, \gamma, \frac{\kappa^2 \phi}{\nu})$$

$$\kappa^2 / \sigma^2 \sim \chi^2(\nu)$$

21

Normal-Inverted Chi (cont.)

- marginal

$$\begin{aligned} h(\mu) &= \\ \int_0^\infty h(\mu, \sigma) d\sigma &\propto \int_0^\infty \sigma^{-\nu-2} \exp\left[-\frac{1}{2\sigma^2} \left(\frac{1}{\phi}(\mu - \gamma)^2 + \kappa^2\right)\right] d\sigma \\ &= \frac{2^{\frac{\nu-1}{2}} \Gamma(\frac{\nu+1}{2})}{\left(\frac{1}{\phi}(\mu - \gamma)^2 + \kappa^2\right)^{\frac{\nu+1}{2}}} \\ &\propto \frac{1}{\left(\frac{1}{\phi}(\mu - \gamma)^2 + \kappa^2\right)^{\frac{\nu+1}{2}}} \\ &\propto \left(\frac{1}{\kappa^2 \phi} (\mu - \gamma)^2 + 1\right)^{-\frac{1}{2}(\nu+1)} \end{aligned}$$

$$\mu \sim t(\nu, \gamma, \frac{\kappa^2 \phi}{\nu})$$

22

Normal-Inverted Chi (cont.)

- moments

$$\mathcal{E}(\mu) = \gamma \quad var(\mu) = \frac{\kappa^2 \phi}{\nu - 2}$$

joint mode(μ) = *marginal mode*(μ) = γ

$$\mathcal{E}(\sigma) \approx \frac{\kappa}{\nu - \frac{3}{2}} \quad var(\sigma) \approx \frac{\kappa^2}{2(\nu - 2)(\nu - \frac{5}{3})}$$

joint mode(σ) = $\frac{\kappa}{\sqrt{\nu + 2}}$ *marginal mode*(σ) = $\frac{\kappa}{\sqrt{\nu + 1}}$

23

Multivariate Normal-Inverted Chi

- notation
- pdf

$$\begin{pmatrix} \boldsymbol{\beta} \\ \sigma \end{pmatrix} \sim MVN\chi^{-1}(\boldsymbol{\gamma}, \boldsymbol{\Phi}, \nu, \kappa^2)$$

$$\begin{aligned} h(\boldsymbol{\beta}, \sigma) &\propto \frac{1}{\sigma^p \sqrt{|\boldsymbol{\Phi}|}} \exp\left(-\frac{1}{2\sigma^2} (\boldsymbol{\beta} - \boldsymbol{\gamma})' \boldsymbol{\Phi}^{-1} (\boldsymbol{\beta} - \boldsymbol{\gamma})\right) \times \\ &\quad \frac{1}{2^{\frac{\nu}{2}-1} \Gamma(\frac{\nu}{2})} \exp\left(-\frac{\kappa^2}{2\sigma^2}\right) (\kappa^2)^{\frac{1}{\nu}} \sigma^{-\nu-1} \\ &= \frac{1}{2^{\frac{\nu}{2}-1} \Gamma(\frac{\nu}{2})} \frac{1}{\sqrt{|\boldsymbol{\Phi}|}} \exp\left[-\frac{1}{2\sigma^2} \left((\boldsymbol{\beta} - \boldsymbol{\gamma})' \boldsymbol{\Phi}^{-1} (\boldsymbol{\beta} - \boldsymbol{\gamma}) + \kappa^2\right)\right] (\kappa^2)^{\frac{1}{\nu}} \sigma^{-\nu-p-1} \\ &\propto \exp\left[-\frac{1}{2\sigma^2} \left((\boldsymbol{\beta} - \boldsymbol{\gamma})' \boldsymbol{\Phi}^{-1} (\boldsymbol{\beta} - \boldsymbol{\gamma}) + \kappa^2\right)\right] \sigma^{-\nu-p-1} \end{aligned}$$

- definition

$$\boldsymbol{\beta} | \sigma \sim MVN(\boldsymbol{\gamma}, \sigma^2 \boldsymbol{\Phi})$$

$$\sigma \sim \chi^{-1}(\nu, \kappa^2)$$

$$\kappa^2 / \sigma^2 \sim \chi^2(\nu)$$

$$\boldsymbol{\beta} \sim t(\nu, \boldsymbol{\gamma}, \frac{\kappa^2}{\nu} \boldsymbol{\Phi})$$

24

Multivariate Normal-Inverted Chi (cont.)

- marginal

$$\begin{aligned}
 h(\boldsymbol{\beta}) &= \int_0^\infty h(\boldsymbol{\beta}, \sigma) d\sigma \\
 &\propto \int_0^\infty \exp \left[-\frac{1}{2\sigma^2} \left((\boldsymbol{\beta} - \boldsymbol{\gamma})' \boldsymbol{\Phi}^{-1} (\boldsymbol{\beta} - \boldsymbol{\gamma}) + \kappa^2 \right) \right] \sigma^{-\nu-p-1} d\sigma \\
 &\propto \left[(\boldsymbol{\beta} - \boldsymbol{\gamma})' \boldsymbol{\Phi}^{-1} (\boldsymbol{\beta} - \boldsymbol{\gamma}) + \kappa^2 \right]^{-\frac{1}{2}(\nu+p)} \\
 &\propto \left[\frac{1}{\nu} (\boldsymbol{\beta} - \boldsymbol{\gamma})' \left(\frac{\kappa^2}{\nu} \boldsymbol{\Phi} \right)^{-1} (\boldsymbol{\beta} - \boldsymbol{\gamma}) + 1 \right]^{-\frac{1}{2}(\nu+p)}
 \end{aligned}$$

$$\boldsymbol{\beta} \sim t(\nu, \boldsymbol{\gamma}, \frac{\kappa^2}{\nu} \boldsymbol{\Phi})$$

25

Beta

- notation $X \sim \text{Beta}(\alpha, \beta)$

- pdf

$$f(x|\alpha, \beta) = \frac{1}{B(\alpha, \beta)} \times x^{\alpha-1} (1-x)^{\beta-1}$$

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$

- moments

$$\mathcal{E}(X) = \frac{\alpha}{\alpha+\beta}, \quad var(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$$

$$mode(X) = \frac{\alpha-1}{\alpha+\beta-2}$$

26

Dirichlet

- notation $X \sim \text{Dir}(\alpha)$

- pdf $0 < X_i < 1, i = 1, 2, \dots, p \quad \sum_{i=1}^p X_i < 1$

$$f(\mathbf{x}|\boldsymbol{\alpha}) = \frac{\Gamma(\sum_{j=1}^{p+1} \alpha_j)}{\prod_{j=1}^{p+1} \Gamma(\alpha_j)} \times \left(\prod_{j=1}^p x_j^{\alpha_j - 1} \right) (1 - \sum_{j=1}^p x_j)^{\alpha_{p+1} - 1}$$

- moments

$$\mathbf{E}(X_i) = \frac{\alpha_i}{\alpha_0}, \quad \text{var}(X_i) = \frac{\alpha_i(\alpha_0 - \alpha_i)}{\alpha_0^2(\alpha_0 + 1)}$$

$$\text{mode}(X_i) = \frac{\alpha_i - 1}{\alpha_0 - 2}$$

$$\text{cov}(X_i, X_j) = -\frac{\alpha_i \alpha_j}{\alpha_0^2(\alpha_0 + 1)}$$

$$\text{jointmode}(\mathbf{X}) = \left\{ \frac{\alpha_i - 1}{\alpha_0 - p - 1} \right\}$$

27

Dirichlet (cont.)

- marginal is beta.

$$X_i \sim B(\alpha_i, \alpha_0), i = 1, 2, \dots, p + 1$$

$$\alpha_0 = \sum_{j=1}^{p+1} \alpha_j$$

28

Wishart

- notation

$$\mathbf{X} \sim W(\nu, \Sigma)$$

X is p x p positive definite matrix.

- pdf

$$f(\mathbf{X} | \nu, \Sigma) = \frac{1}{2^{\frac{1}{2}(\nu p)} \pi^{\frac{1}{4}p(p-1)} |\Sigma|^{\frac{\nu}{2}} \prod_{i=1}^p \Gamma(\frac{1}{2}(\nu + 1 - i))} \times$$

$$|\mathbf{X}|^{\frac{1}{2}(\nu - p - 1)} \exp\left(-\frac{1}{2}\text{tr}(\Sigma^{-1} \mathbf{X})\right)$$

- moment

$$\mathcal{E}(\mathbf{X}) = \nu \Sigma$$

$$\text{var}(x_{ij}) = \nu(\sigma_{ij}^2 + \sigma_{ii}\sigma_{jj})$$

$$\text{mode}(\mathbf{X}) = (\nu - p - 1)\Sigma$$

$$\text{cov}(x_{ij}, x_{kl}) = \nu(\sigma_{ik}\sigma_{jl} + \sigma_{il}\sigma_{jk})$$

29

Inverted Wishart

- notation

$$\mathbf{X} \sim W^{-1}(\nu, \Sigma)$$

$$\mathbf{X}^{-1} \sim W(\nu, \Sigma^{-1})$$

- pdf

$$f(\mathbf{X} | \nu, \Sigma) = \frac{|\Sigma|^{\frac{1}{2}\nu}}{2^{\frac{1}{2}(\nu p)} \pi^{\frac{1}{4}p(p-1)} \prod_{i=1}^p \Gamma(\frac{1}{2}(\nu + 1 - i))} \times$$

$$\frac{1}{|\mathbf{X}|^{\frac{1}{2}(\nu + p + 1)}} \exp\left(-\frac{1}{2}\text{tr}(\Sigma \mathbf{X}^{-1})\right)$$

- moment

$$\mathcal{E}(\mathbf{X}) = \frac{1}{\nu - p - 1} \Sigma$$

$$\text{var}(x_{ii}) = \frac{2(\sigma_{ii})^2}{(\nu - p - 1)(\nu - p - 3)}$$

$$\text{mode}(\mathbf{X}) = \frac{1}{\nu + p + 1} \Sigma$$

30